

A CHARACTERIZATION OF LINEARLY SEMISIMPLE GROUPS

AMELIA ÁLVAREZ, CARLOS SANCHO, AND PEDRO SANCHO

ABSTRACT. Let $G = \text{Spec } A$ be an affine K -group scheme and $\tilde{A} = \{w \in A^* : \dim_K A^* \cdot w \cdot A^* < \infty\}$. Let $\langle -, - \rangle : A^* \times \tilde{A} \rightarrow K$, $\langle w, \tilde{w} \rangle := \text{tr}(w\tilde{w})$, be the trace form. We prove that G is linearly reductive if and only if the trace form is non-degenerate on A^* .

1. INTRODUCTION

Let K be a field and let $\bar{K}$ be its algebraic closure. A finite K -algebra (associative with unit) B is separable (i.e., $B \otimes_K \bar{K} = \prod_i M_{n_i}(\bar{K})$) and $\text{g.c.d.}\{n_i, \text{char } K\} = 1$, for every i , if and only if the trace form of B is non-degenerate.

Let G be a finite group of order n and let KG be the enveloping algebra of G . G is linearly semisimple if and only if KG is a separable K -algebra. By Maschke's theorem G is linearly semisimple if and only if $\text{g.c.d.}\{n, \text{char } K\} = 1$. It holds that G is linearly semisimple if and only if the trace form of KG is non-degenerate. This theorem has been generalized to finite-dimensional Hopf algebras and Frobenius algebras (see [LS] and [M]).

Now let $G = \text{Spec } A$ be an affine K -group scheme. Let $\tilde{A} = \{w \in A^* : \dim_K A^* \cdot w \cdot A^* < \infty\} \subseteq A^*$, which is a bilateral ideal of A^* . Let $\text{tr} : \tilde{A} \rightarrow K$ defined by $\text{tr}(\tilde{w}) = \text{trace of the endomorphism of } \tilde{A}, \tilde{w}' \mapsto \tilde{w}\tilde{w}'$, and $\langle -, - \rangle : A^* \times \tilde{A} \rightarrow K$, $\langle w, \tilde{w} \rangle := \text{tr}(w\tilde{w})$, the associated trace form.

In this paper we prove that G is linearly semisimple if and only if the trace form is non-degenerate on A^* , that is, $\varphi : A^* \rightarrow \tilde{A}^*$, $\varphi(w) := \langle w, - \rangle$ is injective. We also prove that G is linearly semisimple if and only if the trace form is non-degenerate on $\tilde{A}$, that is, $\phi : \tilde{A} \rightarrow \tilde{A}^*$, $\phi(\tilde{w}) := \langle -, \tilde{w} \rangle$ is injective, and $\tilde{A}$ is dense in A^* .

$G = \text{Spec } A$ is linearly semisimple if and only if there exists $w_G \in A^{*G}$ such that $w_G(1) = 1$ (see [Sw]). Let $*$: $A \rightarrow A$, $a \mapsto a^*$ be the morphism induced on A by the morphism $G \rightarrow G$, $g \mapsto g^{-1}$ and let $F : A \rightarrow A^*$, $F(a) := w_G(a^* \cdot -)$ be the “Fourier transform”. In this situation we prove that $\phi : \tilde{A} \rightarrow \tilde{A}^*$ values bijectively in A and that the inverse morphism $\phi^{-1} : A \rightarrow \tilde{A}$ coincides with the Fourier transform.

2. PRELIMINARY RESULTS

If G is a smooth algebraic group over an algebraically closed field one may only consider the rational points of G in order to resolve many questions in the theory of linear representations of G . In general, for any K -group, one may regard the functor of points of the group and its linear representations and in this way K -groups and their linear representations may be handled as mere sets, as it is well known.

Let R be a commutative ring (associative with unit). All functors considered in this paper are covariant functors over the category of commutative R -algebras

(with unit). Given an R -module M , the functor $\mathcal{M}$ defined by $\mathcal{M}(S) := M \otimes_R S$ is called a quasi-coherent R -module. The functors $M \rightsquigarrow \mathcal{M}$, $\mathcal{M} \rightsquigarrow \mathcal{M}(R)$ establish an equivalence between the category of R -modules and the category of quasi-coherent R -modules ([A1, 1.12]). In particular, $\text{Hom}_R(\mathcal{M}, \mathcal{M}') = \text{Hom}_R(M, M')$.

If $\mathbb{M}$ and $\mathbb{N}$ are functors of $\mathcal{R}$ -modules, we will denote by $\mathbb{H}om_{\mathcal{R}}(\mathbb{M}, \mathbb{N})$ the functor of $\mathcal{R}$ -modules

$$\mathbb{H}om_{\mathcal{R}}(\mathbb{M}, \mathbb{N})(S) := \text{Hom}_S(\mathbb{M}|_S, \mathbb{N}|_S),$$

where $\mathbb{M}|_S$ is the functor $\mathbb{M}$ restricted to the category of commutative S -algebras. We denote $\mathbb{M}^* = \mathbb{H}om_{\mathcal{R}}(\mathbb{M}, \mathcal{R})$ and given an R -module M , then

$$\mathcal{M}^*(S) = \text{Hom}_S(M \otimes_R S, S) = \text{Hom}_R(M, S).$$

We will say that $\mathcal{M}^*$ is an R -module scheme and it holds that $\mathcal{M}^{**} = \mathcal{M}$ ([A1, 1.10]).

Remark 2.1. *Every morphism $M^* \rightarrow M'^*$ considered in this paper will be a functorial morphism, that is, we will actually have a morphism $\mathcal{M}^* \rightarrow \mathcal{M}'^*$. Observe that the dual morphism is a morphism $\mathcal{M}' \rightarrow \mathcal{M}$, which is induced by a morphism $M' \rightarrow M$. Then $M^* \rightarrow M'^*$ is the dual morphism of $M' \rightarrow M$.*

Let $G = \text{Spec } A$ be an affine R -group and let G^* be the functor of points of G , that is, $G^*(S) = \text{Hom}_{R\text{-alg}}(A, S)$. $\mathcal{A}^*$ is an R -algebra scheme, that is, besides from being an R -module scheme it is a functor of algebras ($\mathcal{A}^*(S)$ is a S -algebra such that $S \subseteq Z(\mathcal{A}^*(S))$). One has a natural morphism of functors of monoids $G^* \rightarrow \mathcal{A}^*$, because $G^*(S) = \text{Hom}_{R\text{-alg}}(A, S) \subset \text{Hom}_R(A, S) = \mathcal{A}^*(S)$. If $\mathbb{M}^*$ is a functor of $\mathcal{R}$ -modules (resp. $\mathcal{R}$ -algebras), any morphism of functors of sets (resp. monoids) $G^* \rightarrow \mathbb{M}^*$ factorizes uniquely through a morphism of functors of modules (resp. algebras) $\mathcal{A}^* \rightarrow \mathbb{M}^*$ ([A1, 5.3]). Therefore, the category of linear (and rational) representations of G is equal to the category of quasi-coherent $\mathcal{A}^*$ -modules ([A1, 5.5]).

Remark 2.2. *For abbreviation, we sometimes use $g \in G$ or $m \in \mathbb{M}$ to denote $g \in G^*(S)$ or $m \in \mathbb{M}(S)$ respectively. Given $m \in \mathbb{M}(S)$ and a morphism of R -algebras $S \rightarrow S'$, we still denote by m its image by the morphism $\mathbb{M}(S) \rightarrow \mathbb{M}(S')$.*

Definition 2.3. *Let $R = K$ be a field. We will say that an affine K -group scheme G is linearly semisimple if it is linearly reductive, that is, if every linear representation of G is completely reducible.*

G is linearly semisimple if and only if $\mathcal{A}^*$ is semisimple, i.e., $\mathcal{A}^* = \prod_i A_i^*$, where A_i^* are simple (and finite) K -algebras ([A1, 6.8]). If K is an algebraically closed field, then $A_i^* = \text{End}_K V_i$, where V_i are the simple G -modules (up to isomorphism), by Wedderburn's theorem.

Let $V_0 = K$ be the trivial representation of G . The unit of $A_0^* = \text{End}_K V_0$, $w_G := 1_0 \in A_0^* \subset \mathcal{A}^*$ is said to be the “invariant integral on G ”. The invariant integral on G is characterized by being G -invariant and normalized, that is, $w_G(1) = 1$ ([A2, 2.8]). Let $\mathbb{N} = \mathbb{M}^*$ be a functor of G -modules (equivalently, a functor of $\mathcal{A}^*$ -modules). The morphism $w_G \cdot : \mathbb{N} \rightarrow \mathbb{N}$, $n \mapsto w_G \cdot n$ is the unique projection of G -modules of $\mathbb{N}$ onto $\mathbb{N}^G$, and $\mathbb{N}^G = w_G \cdot \mathbb{N}$ (see [A2, 2.3, 3.3]).

3. TRACE FORM

Let us assume that $R = K$ is a field.

Let $\mathcal{A}^*$ be a K -algebra scheme and let $\tilde{A} = \{w \in A^* : \dim_K A^* \cdot w < \infty, \dim_K w \cdot A^* < \infty\}$. It holds that $\tilde{A}$ is a bilateral ideal of A^* .

Remark 3.1. *Functorially, $\tilde{A}$ is the maximal K -quasi-coherent bilateral ideal $\mathcal{I} \subset \mathcal{A}^*$: Given $w \in \tilde{A}$, consider the morphism of $\mathcal{A}^*$ -modules $f: \mathcal{A}^* \rightarrow \mathcal{A}^*$, $f(w') := w' \cdot w$. Let $\mathcal{B}^* = \text{Im } f = \mathcal{A}^* / \ker f$. Observe that $B^* = A^* \cdot w$, then $\dim_K B^* < \infty$ and $\mathcal{B}^*$ is a K -quasi-coherent module. Since $B^* \subset \tilde{A}$, then $\mathcal{B}^* \subset \tilde{A}$ and $\tilde{A}$ is a left $\mathcal{A}^*$ -submodule of $\mathcal{A}^*$. Likewise, $\tilde{A}$ is a right $\mathcal{A}^*$ -submodule of $\mathcal{A}^*$. Hence, $\tilde{A} \subseteq \mathcal{I}$. If $\mathcal{M}$ is a K -quasi-coherent left $\mathcal{A}^*$ -module (resp. right $\mathcal{A}^*$ -module) and $m \in M$, then $\dim_K A^* \cdot m < \infty$ (resp. $\dim_K m \cdot A^* < \infty$) by [A1, 4.7]. Now it follows easily that $\mathcal{I} = \tilde{A}$.*

Definition 3.2. *Let $\mathcal{A}^*$ be a K -algebra scheme. Let $\text{tr}: \tilde{A} \rightarrow K$ be defined by $\text{tr}(\tilde{w}) := \text{trace of the endomorphism of } \tilde{A}, \tilde{w}' \mapsto \tilde{w}\tilde{w}'$. We will say that*

$$\langle -, - \rangle : \mathcal{A}^* \times \tilde{A} \rightarrow K, \quad \langle w, \tilde{w} \rangle := \text{tr}(w\tilde{w})$$

is the trace form of $\mathcal{A}^$.*

We have the associated “polarities” to the trace form

$$\varphi: \mathcal{A}^* \rightarrow \tilde{A}^*, \quad \varphi(w) := \langle w, - \rangle, \quad \phi: \tilde{A} \rightarrow \mathcal{A}^{**} = \mathcal{A}, \quad \phi(\tilde{w}) := \langle -, \tilde{w} \rangle.$$

The dual morphism $\varphi^*: \mathcal{A}^* \rightarrow \tilde{A}^*$ of φ is equal to ϕ :

$$\varphi^*(w)(\tilde{w}) = w(\varphi(\tilde{w})) = \phi(\tilde{w})(w) = \langle w, \tilde{w} \rangle = \varphi(w)(\tilde{w})$$

Likewise, the dual morphism of ϕ is φ .

Definition 3.3. *Let $G = \text{Spec } A$ an affine K -group scheme. We will say that the trace form of $\mathcal{A}^*$ is the trace form of G .*

Let C be a K -algebra ($K \subseteq Z(C)$). If M is a left C -module (resp. a right C -module), then $M^* = \text{Hom}_K(M, K)$ is a right C -module (resp. a left C -module): $(w \cdot c)(m) := w(c \cdot m)$ (resp. $(c \cdot w)(m) := w(m \cdot c)$) for all $w \in M^*$, $m \in M$ and $c \in C$.

The morphism $\varphi: \mathcal{A}^* \rightarrow \tilde{A}^*$ is a morphism of left and right $\mathcal{A}^*$ -modules:

$$\begin{aligned} \varphi(w_1 w_2)(\tilde{w}) &= \text{tr}(w_1 w_2 \tilde{w}) = \text{tr}(w_2 \tilde{w} w_1) = \varphi(w_2)(\tilde{w} w_1) = (w_1 \varphi(w_2))(\tilde{w}) \\ \varphi(w_1 w_2)(\tilde{w}) &= \text{tr}(w_1 w_2 \tilde{w}) = \varphi(w_1)(w_2 \tilde{w}) = (\varphi(w_1) w_2)(\tilde{w}) \end{aligned}$$

for all $w_1, w_2 \in \mathcal{A}^*$ and $\tilde{w} \in \tilde{A}$. Likewise, ϕ is a morphism of left and right $\mathcal{A}^*$ -modules.

If $A^* = \prod_i A_i^*$, where A_i^* are finite K -algebras, then $\tilde{A} = \oplus_i A_i^*$. In this case we have the morphism $\phi: \tilde{A} = \oplus_i A_i^* \rightarrow \oplus_i A_i = A$. Given $1_i := (0, \dots, \overset{i}{1}, \dots, 0) \in \tilde{A}$, then $\phi(1_i) = \text{tr}_{A_i^*}$. Obviously, $\tilde{A}$ is an A^* -module generated by $\{1_i\}_i$, and

$$\langle 1_i, 1_j \rangle = \text{tr}(1_i \cdot 1_j) = \begin{cases} \dim_K A_i^* & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

Example 3.4. Let $G_m = \text{Spec } \mathbb{C}[x, 1/x]$ be the multiplicative group. It holds that $\mathbb{C}[x, 1/x]^* = \prod_{\mathbb{Z}} \mathbb{C}$, $\widehat{\mathbb{C}[x, 1/x]} = \oplus_{\mathbb{Z}} \mathbb{C}$ and

$$\phi: \widehat{\mathbb{C}[x, 1/x]} = \oplus_{\mathbb{Z}} \mathbb{C} \rightarrow \oplus_{n \in \mathbb{Z}} \mathbb{C} \cdot x^n = \mathbb{C}[x, 1/x], \quad \phi((a_n)_{n \in \mathbb{Z}}) = \sum_n a_n x^n.$$

Let $G_a = \text{Spec } \mathbb{C}[x]$ be the additive group. It holds that $\mathbb{C}[x]^* = \mathbb{C}[[z]]$ ($z^n(x^m) = \delta_{nm} \cdot n!$) and $\widehat{\mathbb{C}[[z]]} = 0$.

Let V be a linear representation of an affine K -group scheme $G = \text{Spec } A$. The associated character $\chi_V \in A$ is defined by $\chi_V(g) = \text{trace of the linear endomorphism } V \rightarrow V, v \mapsto g \cdot v$, for every $g \in G$ and $v \in V$.

Let $G = \text{Spec } A$ be a linearly semisimple affine K -group scheme. One has that $A^* = \prod_i A_i^*$, where A_i^* are finite simple algebras. Assume for simplicity that K is an algebraically closed field, then $A_i^* = \text{End}_K(V_i)$. Observe that $\text{tr}_{A_i^*} = n_i \cdot \chi_{V_i}$, where $n_i = \dim_K V_i$.

Let $V_0 = K$ be the trivial representation of G and let $w_G := 1_0 \in A^*$ be the “invariant integral on G ”.

Given $a \in A$, then $w_G \cdot a \in K = A^G$. Hence $w_G \cdot a = (w_G \cdot a)(1) = a(w_G) = w_G(a)$.

One has that $w_G \cdot A_j = 1_0 \cdot A_j = 0$, if $j \neq 0$ and $w_G \cdot a_0 = a_0$ for all $a_0 \in A_0$. Hence, $w_G \cdot \chi_{V_j} = 0$ if V_j is not the trivial representation, and $w_G \cdot \chi_{V_0} = \chi_{V_0} = 1$. Moreover, since $\chi_{V \oplus V'} = \chi_V + \chi_{V'}$, one has

$$(1) \quad w_G(\chi_V) = w_G \cdot \chi_V = \dim_K V^G.$$

4. CHARACTERIZATION OF LINEARLY SEMISIMPLE GROUPS

Let $G = \text{Spec } A$ be an affine K -group scheme.

Let $*$: $A \rightarrow A$, $a \mapsto a^*$ be the morphism induced by the morphism $G \rightarrow G$, $g \mapsto g^{-1}$. In the following proof, if V_i is a linear representation of G , we will consider V_i^* as a left G -module by $(g * w)(v) = w(g^{-1} \cdot v)$. One has that $\chi_{V_i^*} = \chi_{V_i}^*$, because the trace of $g^{-1} \in G$ operating on V_i is equal to the trace of g operating on V_i^* (which operates by the transposed inverse of g).

Theorem 4.1. Let $G = \text{Spec } A$ be a linearly semisimple affine K -group scheme and let $w_G \in A^*$ be its invariant integral. Let

$$F: A \rightarrow A^*, \quad F(a) := w_G(a^* \cdot -)$$

be the “Fourier Transform” (where $w_G(a^* \cdot -)(a') := w_G(a^* \cdot a')$). Then $F(A) = \tilde{A}$ and F is equal to the inverse morphism of the polarity ϕ associated to the trace form of G .

Proof. First let us prove that $F: \mathcal{A} \rightarrow \mathcal{A}^*$, $F(a) := w_G(a^* \cdot -)$ is a morphism of left G -modules. For every point $g \in G$,

$$\begin{aligned} w_G((g \cdot a)^* \cdot -) &\stackrel{*}{=} w_G((a^* \cdot g^{-1}) \cdot -) \stackrel{**}{=} w_G(((a^* \cdot g^{-1}) \cdot -) \cdot g) = w_G(a^* \cdot (- \cdot g)) \\ &= g \cdot (w_G(a^* \cdot -)) \end{aligned}$$

where $*$ is due to $(g \cdot a)^*(g') = a(g'^{-1} \cdot g) = a((g^{-1} \cdot g')^{-1}) = (a^* \cdot g^{-1})(g')$, and $**$ is due to $g \cdot w_G = w_G$. Likewise, F is a morphism of right G -modules. Then F is a morphism of left and right A^* -modules and $F(A) \subseteq \tilde{A}$.

Assume K is an algebraically closed field. Then $A^* = \prod_i A_i^*$, where $A_i^* = \text{End}_K(V_i) = M_{n_i}(K)$. Observe that $M_{n_i}(K) \simeq M_{n_i}(K)^*$ as left and right $M_{n_i}(K)$ -modules. Then $\tilde{A} = \oplus_i M_{n_i}(K)^*$ and $A = \oplus_i M_{n_i}(K)$ are isomorphic as left and right A^* -modules. Let us see that $F \circ \phi = z \cdot$, where $z = (z_i) \in \prod_i K = Z(A^*)$:

$$F \circ \phi \in \text{Hom}_{A^* \otimes A^*}(\tilde{A}, \tilde{A}) = \text{Hom}_{A^* \otimes A^*}(A, A) \subseteq \text{Hom}_{A^* \otimes A^*}(A^*, A^*) = Z(A^*).$$

As $F(\text{tr}_{A_i^*}) = F(\phi(1_i)) = z \cdot 1_i = z_i \cdot 1_i$, then $F(\chi_{V_i}) = (z_i/n_i) \cdot 1_i$. Observe that

$$F(\chi_{V_i})(\chi_{V_j}) = w_G(\chi_{V_i^* \otimes V_j}) = w_G \cdot \chi_{V_i^* \otimes V_j} = \dim_K \text{Hom}_G(V_i, V_j) = \delta_{ij}.$$

Hence, $1 = F(\chi_{V_i})(\chi_{V_i}) = ((z_i/n_i) \cdot 1_i)(\chi_{V_i}) = (z_i/n_i) \cdot n_i = z_i$ and $F \circ \phi = \text{Id}$. Then ϕ is an injective morphism. Since $\phi(A_i^*) \subseteq A_i$, then $\phi(A_i^*) = A_i$ and $\phi: \tilde{A} = \oplus_i A_i^* \rightarrow \oplus_i A_i = A$ is an isomorphism. As a consequence, $\phi \circ F = \text{Id}$. $\square$

Theorem 4.2. *Let $G = \text{Spec } A$ be an affine K -group scheme. G is linearly semisimple if and only if the polarity $\phi: \tilde{A} \rightarrow A$ associated to the trace form of G is an isomorphism.*

Proof. Let us assume that $\phi: \tilde{A} \simeq A$ is a left and right A^* -module isomorphism. Observe that $1 \in A$ is left and right G -invariant. Let $w_G = \phi^{-1}(1)$. Given $w \in A^*$, $w \cdot w_G = w(1)w_G = w_G \cdot w$. Then it is easy to check that $\text{tr}(w_G) = w_G(1)$ and in general $\langle w_G, w' \rangle = w'(1)w_G(1)$. As $\langle w_G, - \rangle = \phi(w_G) \neq 0$, therefore $w_G(1) \neq 0$. Then, $w' = w_G/w_G(1)$ is a normalized invariant integral on G and G is linearly semisimple.

Reciprocally, if G is linearly semisimple, by the previous theorem ϕ is an isomorphism, whose inverse morphism is the Fourier transform. $\square$

Definition 4.3. *We will say that the trace form of $G = \text{Spec } A$ is non-degenerate on A^* if the associated polarity, $\varphi: A^* \rightarrow \tilde{A}^*$, $\varphi(w) := \langle w, - \rangle$, is an injective morphism.*

Theorem 4.4. *Let $G = \text{Spec } A$ be an affine K -group scheme. G is linearly semisimple if and only if its trace form is non-degenerate on A^* .*

Proof. If $\varphi: A^* \rightarrow \tilde{A}^*$ is injective, then dually $\phi: \tilde{A} \rightarrow A$ is surjective. Let us denote by $i: \tilde{A} \rightarrow A^*$ the natural inclusion. From the commutative diagram

$$\begin{array}{ccc} \tilde{A} & \xrightarrow{\phi} & A \\ \downarrow i & & \downarrow i^* \\ A^* & \xrightarrow{\varphi} & \tilde{A}^* \end{array}$$

it follows that $\phi: \tilde{A} \rightarrow A$ is an isomorphism and we have that G is linearly semisimple.

Reciprocally, if G is linearly semisimple, by the previous theorem $\phi: \tilde{A} \rightarrow A$ is an isomorphism, then dually $\varphi: A^* \rightarrow \tilde{A}^*$ is an isomorphism. $\square$

Given a vector space V , on V^* we will consider the topology whose closed sets are $\{V^0 = (V/V')^*\}_{V' \subseteq V}$. Given a vector subspace $W \subseteq V^*$ we denote $W^0 = \{v \in V: v(W) = 0\}$. The closure of W in V^* coincides with W^0 . (Functorially, the closure of $\mathcal{W}$ in $\mathcal{V}^*$ is the minimum K -scheme of submodules of $\mathcal{V}^*$ which contains $\mathcal{W}$). The closure of $\tilde{A}$ in A^* is a bilateral ideal.

Definition 4.5. We will say that the trace form of $G = \operatorname{Spec} A$ is non-degenerate on $\tilde{A}$ if its associated polarity $\phi: \tilde{A} \rightarrow \tilde{A}^*$, $\phi(\tilde{w}) = \langle -, \tilde{w} \rangle$ is an injective morphism.

Theorem 4.6. Let $G = \operatorname{Spec} A$ be an affine K -group scheme. G is a linearly semisimple group if and only if $\tilde{A}$ is dense in A^* and its trace form is non-degenerate on $\tilde{A}$.

Proof. Let us assume that the trace form of $G = \operatorname{Spec} A$ is non-degenerate on $\tilde{A}$. Then, the morphism $\phi: \tilde{A} \rightarrow A$ is injective, hence dually the morphism $\varphi: A^* \rightarrow \tilde{A}^*$ is surjective. $\ker \varphi$ is a closed set of A^* and $\ker \varphi \cap \tilde{A} = \ker \phi = 0$. Taking closure we obtain that $\ker \varphi \cap A^* = 0$, therefore $\ker \varphi = 0$.

In conclusion, $\varphi: A^* \rightarrow \tilde{A}^*$ is an isomorphism, then dually $\phi: \tilde{A} \rightarrow A$ is an isomorphism and we obtain that G is linearly semisimple. $\square$

5. PONTRJAGIN'S DUALITY

Now our aim is to write Theorem 4.6 in terms of the classical Pontrjagin's duality, which establishes the duality between compact commutative groups and discrete commutative groups.

$\tilde{A}$ is dense in A^* if and only if for every epimorphism of algebra schemes $\pi: \mathcal{A}^* \rightarrow \mathcal{B}^*$, where $\dim_K \mathcal{B}^* < \infty$, it holds that $\pi(\tilde{A}) = \mathcal{B}^*$. Since $\mathcal{B}^*$ injects into $\operatorname{End}_K(\mathcal{B}^*)$, we obtain the following proposition.

Proposition 5.1. Let $G = \operatorname{Spec} A$ be an affine K -group scheme. $\tilde{A}$ is dense in A^* if and only if for every linear representation, $G \rightarrow \operatorname{End}_K(V)$, where $\dim_K V < \infty$, the image of the induced morphism $A^* \rightarrow \operatorname{End}_K(V)$ coincides with the image of $\tilde{A}$.

Let us assume that $\tilde{A}$ is dense in A^* . We want to determine the maximal bilateral ideals of $\tilde{A}$.

Let V be a G -module. For every $v \in V$, it holds that $v \in \tilde{A} \cdot v$: the G -module $\langle v \rangle$ generated by v is finite-dimensional, then there exists $w \in \tilde{A}$ such that the morphism $w: \langle v \rangle \rightarrow \langle v \rangle$, $v' \mapsto w \cdot v'$ is the identity morphism. Then $v = w \cdot v \in \tilde{A} \cdot v$.

In particular, for every $\tilde{w} \in \tilde{A}$ it holds that $\tilde{w} \in \tilde{A} \cdot \tilde{w}$.

Let $I \subseteq \tilde{A}$ be an ideal and $w \in A^*$, then $w \cdot I \subseteq I$: $w \cdot I \subseteq \tilde{A} \cdot (w \cdot I) \subseteq \tilde{A} \cdot I \subseteq I$. Then, every bilateral ideal $I \subset \tilde{A}$ is a left and right A^* -submodule, then $\mathcal{I}$ is a left and right $\mathcal{A}^*$ -module, by [A1, 4.6]. If $I \subsetneq \tilde{A}$ is a maximal bilateral ideal, then $\dim_K \tilde{A}/I < \infty$, because if $\mathcal{M}$ is a left and right quasi-coherent $\mathcal{A}^*$ -module and $m \in \mathcal{M}$, then the left and right quasi-coherent $\mathcal{A}^*$ -module generated by m is finite dimensional, by [A1, 4.7].

Let $I \subset \tilde{A}$ be a bilateral ideal of finite codimension and $\tilde{w} \in \tilde{A}$ such that $\tilde{w}: \tilde{A}/I \rightarrow \tilde{A}/I$ is the identity morphism. Then, $\tilde{w} \cdot \bar{v} = \bar{v}$, for every $\bar{v} \in \tilde{A}/I$, that is, $\tilde{w}$ is a left unit of $\tilde{A}/I$. Likewise, there exists a right unit in $\tilde{A}/I$, then there exists a unit $1 \in \tilde{A}/I$. The morphism $\pi: \mathcal{A}^* \rightarrow \tilde{\mathcal{A}}/\mathcal{I}$, $\pi(w) = w \cdot 1$ is a surjective morphism of $\mathcal{K}$ -algebra schemes. If $\mathcal{J}^* = \ker \pi$, then $A^*/\mathcal{J}^* = \tilde{A}/I$. Hence, the algebra scheme closure of $\tilde{\mathcal{A}}$ is $\mathcal{A}^*$: If $\mathcal{B}^*$ is an algebra scheme, then $\mathcal{B}^* = \varprojlim_i \mathcal{B}_i^*$ is

an inverse limit of algebra schemes $\mathcal{B}_i^*$, where $\dim_K \mathcal{B}_i < \infty$, by [A1, 4.12]. Then

$$\begin{aligned} \operatorname{Hom}_{\mathcal{K}\text{-alg}}(\tilde{\mathcal{A}}, \mathcal{B}^*) &= \varprojlim_i \operatorname{Hom}_{\mathcal{K}\text{-alg}}(\tilde{\mathcal{A}}, \mathcal{B}_i^*) = \varprojlim_i \operatorname{Hom}_{\mathcal{K}\text{-alg}}(\mathcal{A}^*, \mathcal{B}_i^*) \\ &= \operatorname{Hom}_{\mathcal{K}\text{-alg}}(\mathcal{A}^*, \mathcal{B}^*). \end{aligned}$$

Let $G = \operatorname{Spec} A$ be an affine K -group scheme and let $\hat{G}$ be the “dual group”, that is, the set of irreducible representations of G , up to isomorphism. If $\tilde{A}$ is dense in A^* , then

$$\begin{aligned} \operatorname{Spec} \tilde{A} &:= \{\text{Maximal bilateral ideals } I \subsetneq \tilde{A}\} \\ &= \{\text{Maximal bilateral ideal schemes } \mathcal{J}^* \subset \mathcal{A}^*\} \stackrel{[A1, 6.11-12]}{=} \hat{G}. \end{aligned}$$

Definition 5.2. *We will say that $\hat{G}$ is discrete if $\tilde{A}$ is dense in A^* and $\tilde{A}$ is separable (that is, $\tilde{A}$ is a direct sum of algebras of matrices by base change to the algebraic closure of K).*

Now we can rewrite Theorem 4.6 as follows.

Theorem 5.3. *Let $G = \operatorname{Spec} A$ be an affine K -group scheme. G is linearly semisimple if and only if $\hat{G}$ is a discrete group.*

Proof. If G is linearly semisimple, then $A^* = \prod_i A_i^*$, where A_i^* is a finite simple algebra for all i . By base change G is linearly semisimple, then A_i^* is a direct sum of algebras of matrices by base change to the algebraic closure of K . Since $\tilde{A} = \oplus_i A_i^*$, then $\tilde{A}$ is dense in A^* and it is separable.

If $\tilde{A}$ is separable then its trace form is non-degenerate. Hence, if $\tilde{A}$ is separable and dense in A^* , then G is linearly semisimple by Theorem 4.6. $\square$

REFERENCES

- [A1] ÁLVAREZ, A., SANCHO, C., SANCHO, P., *Algebra schemes and their representations*, J. Algebra **296/1** (2006) 110-144.
- [A2] ÁLVAREZ, A., SANCHO, C., SANCHO, P., *Reynolds operator*, arXiv:math/0611311v3 [math.AG] (2008).
- [LS] LARSON, R.G., SWEEDLER, M.E., *An associative orthogonal bilinear form for Hopf algebras*, Am. J. Math. **91** (1969) 75-93.
- [M] MURRAY, W., *Bilinear forms on Frobenius algebras*, J. Algebra **293** (2005) 89-101.
- [S] SANCHO DE SALAS, C., *Grupos algebraicos y teoría de invariantes*, Sociedad Matemática Mexicana, México, 2001.
- [Sw] SWEEDLER, M.E., *Hopf Algebras*, Mathematics Lecture Note Series 44, Addison-Wesley, New York, 1969.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, AVENIDA DE ELVAS S/N, 06071 BADAJOZ, SPAIN

E-mail address: `aalarma@unex.es`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE SALAMANCA, PLAZA DE LA MERCED 1-4, 37008 SALAMANCA, SPAIN

E-mail address: `mplu@usal.es`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, AVENIDA DE ELVAS S/N, 06071 BADAJOZ, SPAIN

E-mail address: `sancho@unex.es`